

Manoj P. Reddy, MD

Sports Medicine & Shoulder Surgery
Orthopedic Associates of Dallas
469.800.7200 office | 469.800.7210 fax
DallasOrtho.com
ReddyOrtho.com

Patella ORIF Protocol

Phase I: 0-2 Weeks
 Knee Brace: Worn at all times except hygiene and Physical therapy. If placed in cast will be changed to brace at 1st visit Weightbearing: WBAT with the knee locked in extension Range of Motion: AROM/AAROM/PROM 0-30 degrees Therapeutic Exercises: Isometric quadriceps/hamstring/adductor/abductor strengthening, Ankle theraband exercises
 Phase II: 2-6 Weeks Knee Brace: Worn with weightbearing activities still locked in full extension- may be removed at night Weightbearing:Range of Motion: AROM/AAROM/PROM - add 15 degrees of flexion each week - Goal is 90 degrees by post-op week 6 Therapeutic Exercises: Isometric quadriceps/hamstring/adductor/abductor strengthening, Ankle theraband exercises, Initiate straight leg raises
 Phase III: 6-10 Weeks Knee Brace: Unlocked – worn with weightbearing activities Weightbearing: Full Range of Motion: AROM/AAROM/PROM – progress to full ROM by post-operative week 10 Therapeutic Exercises: Isometric quadriceps/hamstring/adductor/abductor strengthening, Ankle theraband exercises, Initiate straight leg raises
Phase IV: 10-12 Weeks • Knee Brace: Discontinue • Weightbearing: Full • Range of Motion: Full • Therapeutic Exercises: Isometric quadriceps/hamstring/adductor/abductor strengthening, Ankle theraband exercises, Initiate straight leg raises, Start stationary bicycle
Phase V: 3-6 Months • Return to full activities as tolerated