

Insurance Verification Form

Fax with copies of insurance card(s), front and back, to Amgen® SupportPlus: 1-877-877-6542



Patient Information New Patient to Prolia® Existing Patient

*Patient Name: _____
 Attach patient demographic sheet **OR** Complete information below:
*Street Address: _____
*City: _____ *State: _____ *ZIP: _____
*Phone: _____
M F *Date of Birth: _____

Fulfillment Method (Select only ONE)

Medical Benefit (Physician Purchase)
 Pharmacy Benefit Out of Network Benefits
 Referral to treating site:
*Enter Site ID: _____ **OR** Complete information below.
*Site Name: _____
*Street Address: _____
*City: _____ *State: _____ *ZIP: _____
*Phone: _____ *Fax: _____
Office Contact: _____
*Site Type: MD Office Hospital Outpatient

Primary Insurance Information

Attach a copy of insurance card, front AND back **OR** provide:
*Insurance Name: _____
*Insurance Phone: _____
Subscriber Name: _____
Subscriber Date of Birth: _____
Subscriber Relationship to Patient: _____
Group #: _____
*Policy #: _____
Medicare Beneficiary Identifier: _____

Secondary Insurance Information (If Applicable)

Attach a copy of insurance card, front AND back **OR** provide:
*Insurance Name: _____
*Is this a Medigap policy? Yes No Not Known
If yes, please indicate plan letter: _____
*Insurance Phone: _____
Subscriber Name: _____
Subscriber Date of Birth: _____
Subscriber Relationship to Patient: _____
Group #: _____
*Policy #: _____

Pharmacy Insurance Information

Attach a copy of insurance card, front AND back **OR** provide:
*Pharmacy Insurance Patient ID #: _____
*Pharmacy Insurance Phone #: _____

*Asterisk fields are required for processing.

Physician Information

*Physician Name: _____
*NPI #: _____ Tax ID #: _____
Specialty: _____
*Enter Site ID: _____ **OR** Complete information below.
*Site NPI #: _____ Site Tax ID #: _____
*Site Name: _____
*Street Address: _____
*City: _____ *State: _____ *ZIP: _____
*Phone: _____ Fax: _____
Office Contact: _____
*Site Type: MD Office Hospital Outpatient

Patient Medical Information†

M81.0 (Age-related osteoporosis without current pathological fracture)
 M80.0 _____ (Age-related osteoporosis with current pathological fracture...) Please provide complete code
 Other (specify ICD Code) _____
Please provide secondary ICD Code, if applicable: _____
Please NOTE: clinical notes and additional documentation are **NOT required** for us to process a patient benefit verification. Review of clinical documentation sent to Amgen SupportPlus could delay our response time back to your office. Please **DO NOT** provide anything beyond the information requested on this benefit verification form.
†The sample diagnosis codes are informational and not intended to be directive or a guarantee of reimbursement and include potential codes that would include FDA approved indications for Prolia®. Other codes may be more appropriate given internal system guidelines, payer requirements, practice patterns, and the services rendered.

Prescription Information

Prolia® 60 mg pre-filled syringe, 60 mg SC every 6 months
Refill: x1
Prescriber Signature: (required for legal prescription triage)
X _____ Date: _____

Injection Date

Patient's Scheduled Injection Date: _____

OPTIONAL: Affordability Screening

To see if the patient is eligible for additional affordability options, please complete the questions below
Residency:
Patient has lived in the U.S. or its territories (American Samoa, Guam, Puerto Rico, or U.S. Virgin Islands): Greater than 6 months
 Less than 6 months
Patient household income: \$ _____ Monthly Annually
(Gross income includes all individuals in the household. This includes wages, Social Security, Social Security disability, unemployment, pensions, and any other income. They may be asked to provide proof of income.)
How many people live in the patient's household (including the patient)?:
 1 2 3 4 Other _____
Household size includes all individuals reported on the patient's U.S. Tax Return. If the patient did not file a tax return please include all individuals that live with them.

By completing and faxing this form, you represent that your patient has requested and authorized the disclosure of their personal health information to Amgen and its agents for Amgen to provide the patient support services described in this paragraph. You represent that you have explained to the patient, and the patient indicated they understand and have consented to, the following: 1) Amgen and its agents will use the patient's name, date of birth, contact information, prescriptions, and other necessary health information listed in this form for reimbursement services related to this prescription, including to verify their insurance benefits, and to contact the patient directly for the administration of these patient support services; 2) Amgen will then disclose the patient's personal information to the insurer(s) listed on this form for the same purposes; 3) the patient can withdraw their consent by contacting Amgen at (866) 264-2778 or visiting www.amgen.com/DataSubjectRights, but if the patient does not agree to, or withdraws consent for, these uses or disclosures, the patient cannot receive these patient support services for this medication which necessarily requires Amgen to process the patient's personal information; 4) the patient can view more details about Amgen's privacy practice at www.amgen.com/privacy.

Indications

Prolia® is indicated for the treatment of postmenopausal women with osteoporosis at high risk for fracture, defined as a history of osteoporotic fracture, or multiple risk factors for fracture; or patients who have failed or are intolerant to other available osteoporosis therapy. In postmenopausal women with osteoporosis, Prolia® reduces the incidence of vertebral, nonvertebral, and hip fractures.

Prolia® is indicated for treatment to increase bone mass in men with osteoporosis at high risk for fracture, defined as a history of osteoporotic fracture, or multiple risk factors for fracture; or patients who have failed or are intolerant to other available osteoporosis therapy.

Important Safety Information

• SEVERE HYPOCALCEMIA IN PATIENTS WITH ADVANCED KIDNEY DISEASE:

Patients with advanced chronic kidney disease are at greater risk of severe hypocalcemia following Prolia administration. Severe hypocalcemia resulting in hospitalization, life-threatening events and fatal cases have been reported. The presence of chronic kidney disease-mineral bone disorder (CKD-MBD) markedly increases the risk of hypocalcemia. Prior to initiating Prolia in patients with advanced chronic kidney disease, evaluate for the presence of CKD-MBD. Treatment with Prolia in these patients should be supervised by a healthcare provider with expertise in the diagnosis and management of CKD-MBD.

- **Contraindications:** Prolia® is contraindicated in patients with hypocalcemia. Pre-existing hypocalcemia must be corrected prior to initiating Prolia®. Prolia® is contraindicated in women who are pregnant and may cause fetal harm. In women of reproductive potential, pregnancy testing should be performed prior to initiating treatment with Prolia®. Prolia® is contraindicated in patients with a history of systemic hypersensitivity to any component of the product. Reactions have included anaphylaxis, facial swelling and urticaria.

- **Severe Hypocalcemia and Mineral Metabolism Changes:** Prolia can cause severe hypocalcemia and fatal cases have been reported. Pre-existing hypocalcemia must be corrected prior to initiating therapy with Prolia. Adequately supplement all patients with calcium and vitamin D.

In patients without advanced chronic kidney disease who are predisposed to hypocalcemia and disturbances of mineral metabolism (e.g. treatment with other calcium-lowering drugs), assess serum calcium and mineral levels (phosphorus and magnesium) 10 to 14 days after Prolia injection.

- **Same Active Ingredient:** Patients receiving Prolia® should not receive other denosumab products concomitantly.
- **Hypersensitivity:** Clinically significant hypersensitivity including anaphylaxis has been reported with Prolia®. Symptoms have included hypotension, dyspnea, throat tightness, facial and upper airway edema, pruritus and urticaria. If an anaphylactic or other clinically significant allergic reaction occurs, initiate appropriate therapy and discontinue further use of Prolia®.
- **Osteonecrosis of the Jaw (ONJ):** ONJ, which can occur spontaneously, is generally associated with tooth extraction and/or local infection with delayed healing, and has been reported in patients receiving Prolia®. An oral exam should be performed by the prescriber prior to initiation of Prolia®. A dental examination with appropriate preventive dentistry is recommended prior to treatment in patients with risk factors for ONJ such as invasive dental procedures, diagnosis of cancer, concomitant therapies (e.g. chemotherapy, corticosteroids, angiogenesis inhibitors), poor oral hygiene, and co-morbid disorders. Good oral hygiene practices should be maintained during treatment with Prolia®. The risk of ONJ may increase with duration of exposure to Prolia®.
For patients requiring invasive dental procedures, clinical judgment should guide the management plan of each patient. Patients who are suspected of having or who develop ONJ should receive care by a dentist or an oral surgeon. Extensive dental surgery to treat ONJ may exacerbate the condition. Discontinuation of Prolia® should be considered based on individual benefit-risk assessment.
- **Atypical Femoral Fractures:** Atypical low-energy, or low trauma fractures of the shaft have been reported in patients receiving Prolia®. Causality has not been established as these fractures also occur in osteoporotic patients who have not been treated with antiresorptive agents.

During Prolia® treatment, patients should be advised to report new or unusual thigh, hip, or groin pain. Any patient who presents with thigh or groin pain should be evaluated to rule out an incomplete femur fracture. Interruption of Prolia® therapy should be considered, pending a risk/benefit assessment, on an individual basis.

- **Multiple Vertebral Fractures (MVF) Following Discontinuation of Prolia® Treatment:** Following discontinuation of Prolia® treatment, fracture risk increases, including the risk of multiple vertebral fractures. New vertebral fractures occurred as early as 7 months (on average 19 months) after the last dose of Prolia®. Prior vertebral fracture was a predictor of multiple vertebral fractures after Prolia® discontinuation. Evaluate an individual's benefit/risk before initiating treatment with Prolia®. If Prolia® treatment is discontinued, patients should be transitioned to an alternative antiresorptive therapy.
- **Serious Infections:** In a clinical trial (N = 7808), serious infections leading to hospitalization were reported more frequently in the Prolia® group than in the placebo group. Serious skin infections, as well as infections of the abdomen, urinary tract and ear, were more frequent in patients treated with Prolia®.
Endocarditis was also reported more frequently in Prolia®-treated patients. The incidence of opportunistic infections and the overall incidence of infections were similar between the treatment groups. Advise patients to seek prompt medical attention if they develop signs or symptoms of severe infection, including cellulitis. Patients on concomitant immunosuppressant agents or with impaired immune systems may be at increased risk for serious infections. In patients who develop serious infections while on Prolia®, prescribers should assess the need for continued Prolia® therapy.
- **Dermatologic Adverse Reactions:** Epidermal and dermal adverse events such as dermatitis, eczema and rashes occurred at a significantly higher rate with Prolia® compared to placebo. Most of these events were not specific to the injection site. Consider discontinuing Prolia® if severe symptoms develop.
- **Musculoskeletal Pain:** Severe and occasionally incapacitating bone, joint, and/or muscle pain has been reported in patients taking Prolia®. Consider discontinuing use if severe symptoms develop.
- **Suppression of Bone Turnover:** Prolia® resulted in significant suppression of bone remodeling as evidenced by markers of bone turnover and bone histomorphometry. The significance of these findings and the effect of long-term treatment are unknown. Monitor patients for consequences, including ONJ, atypical fractures, and delayed fracture healing.
- **Adverse Reactions:** The most common adverse reactions (>5% and more common than placebo) in women with postmenopausal osteoporosis are back pain, pain in extremity, musculoskeletal pain, hypercholesterolemia, and cystitis. The most common adverse reactions (> 5% and more common than placebo) in men with osteoporosis are back pain, arthralgia, and nasopharyngitis. Pancreatitis has been reported with Prolia®.
In women with postmenopausal osteoporosis, the overall incidence of new malignancies was 4.3% in the placebo group and 4.8% in the Prolia® group. In men with osteoporosis, new malignancies were reported in no patients in the placebo group and 4 (3.3%) patients in the Prolia® group. A causal relationship to drug exposure has not been established.
Denosumab is a human monoclonal antibody. As with all therapeutic proteins, there is potential for immunogenicity.



Amgen Inc.
One Amgen Center Drive
Thousand Oaks, CA 91320-1799
www.amgen.com

Please see Prolia® full Prescribing Information, including Medication Guide.

© 2022-2025 Amgen Inc. All rights reserved. USA-162-83767 11/25