PACIFIC INFUSION CENTER
Friendly • Experienced • Caring

1260 15th Street, Suite 1400
Santa Monica, CA 90404
www.pacificinfusion.com
(310) 297-9269 office
(310) 297-9222 fax

Patient Name: _____ **Phone:**_____

Physician use only: **DOB:** _____**Patient Weight (in kg):** _____

SIG:

- ☐ **Actemra** _____ mg per kg IV every 4 weeks
- ☐ **Benlysta** 10 mgs per kg IV on weeks 0, 2, 4 and then every 4 weeks
- ☐ **Cimzia** _____mg SC on weeks 0, 2, 4 and then every 4 weeks
- ☐ **Cosentyx** 6 mg on week 0 as a loading dose then 1.75 mg per kg every 4 weeks
- ☐ **Cosentyx** 1.75 mg every 4 weeks (without loading dose)
- ☐ **Entyvio** 300 mgs IV @ weeks 0, 2, 6 and then every ___weeks
- ☐ **Evenity** 210 mg SC every month for 12 months
- ☐ **Gazyva** 1000 mg IV @ weeks 0, 2, 24, 26 then every 6 weeks
- ☐ **Ilumya** 100 mgs SC @ weeks 0, 4 and then 100 mg SC every 12 weeks
- ☐ **Imaavy** 30 mgs IV @ weeks 0 and then 15/mg/kg IV every 2 weeks (generalized myasthenia gravis)
- ☐ **Imaavy** 30 mgs IV every 4 weeks (Warm autoimmune hemolytic anemia)
- ☐ **Kisunla** 700 mgs IV @ weeks 0, 4 and 8 then 1400 mg every 4 weeks
- ☐ **Leqembi** 10 mgs per kg IV every 2 weeks
- ☐ **Leqvio** 284 mgs SC on week 0, then at 3 months then every 6 months
- ☐ **Ocrevus** 300 mgs IV @ weeks 0, 2 and then 600mg IV every 6 months
- ☐ **Ocrevus Zunovo** 23mL SC every 6 months
- ☐ **Omvoh** 300 mgs IV @ weeks 0, 4 and 8
- ☐ **Orencia** _____mgs IV on weeks 0, 2, 4 and then every 4 weeks
- ☐ **Prolia** 60 mg SC every 6 months
- ☐ **Remicade** _____mgs per kg IV @ weeks 0, 2, 6 and then every ___weeks
- ☐ **Renflexis** _____mgs per kg IV @ weeks 0, 2, 6 and then every ___weeks
- ☐ **Rituxan** _____ mgs IV every _____
- ☐ **Saphenelo** 300 mgs IV every 4 weeks
- ☐ **Simponi Aria** 2 mg per kg IV at weeks 0, 4 and then every 8 weeks
- ☐ **Skyrizi** 600 mgs IV @ weeks 0, 4, 8 (Crohn's Disease)
- ☐ **Skyrizi** 1200 mgs IV @ weeks 0, 4, 8 (Ulcerative Colitis)
- ☐ **Stoboclo** 60 mg SC every 6 months
- ☐ **Stelara** _____mgs IV @ weeks 0, 4 and then _____ mg IV every ____ weeks
- ☐ **Tepezza** 10 mgs per kg IV @ weeks 0 an then 20 mg per kg IV every 3 weeks
- ☐ **Tremfya** 200 mgs IV @ weeks 0, 4, 8 (Induction dose for Ulcerative Colitis and Crohn's Disease)
- ☐ **Uplizna** 300 mgs @ weeks 0, 2 and then 300mg every 6 months from week 0
- ☐ **Other:** _____ ☐ IM or ☐ IV or ☐ SC q _____
- ☐ **Do not substitute**

Premedication?: ☐ **Yes** ☐ **No**

- ☐ **Acetaminophen** _____mg PO
- ☐ **Diphenhydramine** 25mg IV
- ☐ **Fexofenadine** 180mg PO ☐ **Screening labs/tests sent to us**
- ☐ **Methyprednisolone** ☐ 40mg IV ☐ 125mg ☐ Other: _____
- ☐ _____

Dx: _____ **ICD-10 code:**_____

MD Signature: _____ **MD Print Name:** _____

Date: _____ **Phone Number:** _____ **Fax Number:** _____

Fax this prescription to our office with facesheet/insurance card/requested labs and/or tests (see www.pacificinfusion.com for comprehensive list) & give copy to patient.