

PRE-OP NURSE

- All piercings removed
- All rings removed
- Lovenox 40 mg SC in thigh – NOT abdomen
- Chlorhexidine wipes to chest/axilla/abdomen/groin
- Mepilex on sacrum

CIRCULATING NURSE

- Warm room before patient enters until all drapes applied
- Check OR table to make sure the back raises
- Underbody warmer (if available) and lower body Bair hugger
- Foley catheter with temp probe – padded under thigh
- SCDs
- Microscope available
- SPY available
- Vioptix monitor plugged in & turned on
- Turn heels every 2 hours to prevent pressure sore
- Check wrists/hands when patient flexed for closure

- Call family to give updates
 - 11 am
 - 4 pm

SURGEON

- Patient positioning
- Remove all arm bands
- Pad all IV lines
- Communicate with anesthesia about intraop goal directed therapy
- Surgical site preparation and draping
- Debrief

ANESTHESIA

- All EKG leads on back
- Pulse oximeter on forehead or ear – NOT finger
- No Arterial line needed
- Prophylactic antibiotics and redosing schedule
- IV and BP cuff on unrestricted limb
- Second IV in foot – remove SCD on that side
- No pressure bag on IVF
- Euvolemia ~ 300 cc per hour
- Endotracheal intubation
- Muscle paralysis
- ESP and TAP blocks (with Exparel) once intubated
- Communicate with surgeon if pressors needed
- Turn head every 2 hours to prevent occipital alopecia

- ERAS Protocol – see additional sheet